

Division of Signal Transduction Therapy

Standard Operating Procedure

Preparation of ABIN3 D210N L228P [1 - 325]

Enzyme description:- ABIN3 D210N L228P [1 – 325]

Clone number:- DU 8825

Source:- Recombinant

Expression system:- *E.coli*

Tag:- N-terminal GST

Purification method:- GSH Sepharose

Calculated molecular mass:-

Monoisotopic 66, 626.64 daltons

Average Mass 66, 669.22 daltons

[cysteines reduced, methionines have not been oxidised]

Theoretical pI:- 6.78

Purity:- >80 %

Enzyme storage buffer:-

50 mM Tris-HCl pH 7.5, 270 mM Sucrose, 150 mM NaCl, 0.1 mM EGTA,
0.1 % 2-mercaptoethanol, 0.02 % Brij-35, 1 mM benzamidine, 0.2 mM PMSF

Storage temperature:- -70 °C

Division of Signal Transduction Therapy

Clone Data Sheet

ABIN3 D210N L228P [1 – 325]

<u>Protein</u>	ABIN3 D210N L228P [1 – 325]
<u>Clone number</u>	DU 8825
<u>Species</u>	Human
<u>Accession number</u>	NM_024873.5
<u>Tags</u>	N-terminal GST
<u>Bacterially expressed protein</u>	<p>MSPILGYWKIKGLVQPTRLLEYLEEKYEEHLYERDEGDKWRNKKFELG LEFPNLPYYIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAIEISMLEGA VLDIRYGVSR IAYS KDFETLKVDFLSKLP EMLKMFEDRLCHKTYLNGDH VTHPDFMLYDALDVVLYMDPMCLDAFPKLVCFKKRIEAIPOIDKYLKSS KYIAWPLQGWQATFGGGDHPKSDLEVL FQG PLGSPGIPGSTRAAAMAH FVQGT SRMIAAES STEHKECAEP STRKNLMNSLEQKIRCLEKQRKELLE VNQOWDQOFRSMKELYERKVAELKTKLDAAERFLSTREKDPHQQRKDD RQREDDRQRDLTRDLQREEKEKERLNEELHELKEENKLLKGKNTLANK EKEHYECEIKRLNKALQDALNIKCSFSEDCLRKSRVEFCHEEMRTEMEV LKQOVQIYEENFKKERSDRERLNQEKEEPQQINETSQSQNLNRLNSQIKA CQMEKEKLEKQLKQMYCPPCNCGLVFHLQDPWVPTGPGAVQKQREHPPD CQWYALDQLPPDVQHKANGLSSVKKVHP</p>
<u>Native sequence</u>	<p>Amino acids M1 – P325 (end) of human ABIN3. Residue M243 of the fusion protein is equivalent to M1 of the native enzyme. The GST tag is located at residues 1 – 220. The protein has a D210N and an L228P mutation. Residue D210 is equivalent to residue N452 of the fusion protein. Residue L228 is equivalent to residue P470 of the fusion protein.</p>
<u>Protease cleavage</u>	PreScission (LEVL FQGP) residues 221 - 228
<u>Cloning sites</u>	Not1 sites of pGEX6P-2

Division of Signal Transduction Therapy

Nucleotide Sequence Of Insert

gcggccgcgATGGCACATTTTGTACAGGGCACATCTAGAATGATTGCCGCAGAA
AGTTCTACGGAGCATAAAGAGTGTGCTGAACCATCAACAAGAAAAGAACTTGATG
AATTCTCTTGAACAAAAGATAAGGTGTTTGGAAAAACAAAGAAAAGAGCTCCTG
GAAGTTAACCAGCAATGGGATCAGCAATTTAGAAGTATGAAAGAGTTATATGAA
AGAAAGGTAGCAGAGCTGAAGACGAAACTGGACGCCGCGGAAAGATTCCCTCAGC
ACGCGGGAGAAGGATCCGCATCAGAGGCAGAGAAAAGGACGACAGGCAGAGAGAG
GACGACAGGCAGCGCGACCTGACCCGGGACCGGCTGCAGCGGGAGGAGAAGGAA
AAGGAACGCCTAAATGAAGAATTACATGAATTGAAAAGAGAATAAACTTTTA
AAGGGAAAAAATACTCTTGCGAACAAGGAAAAGGAACATTACGAATGTGAAATA
AAACGCCTCAATAAGGCTCTTCAGGATGCCTTGAATATCAAGTGTTCATTTTCC
GAGGACTGTTTGAGGAAGTCTCGAGTGGAATTCTGCCATGAGGAGATGAGAACA
GAAATGGAAGTTCTGAAGCAGCAGGTGCAAATATACGAAGAAAACCTCAAAAAG
GAACGATCGGATCGAGAGAGACTTAATCAAGAGAAAAGAGGAGCCACAGCAAATT
AATGAAACTTCCCAATCCCAGTTGAACAGGCTGAATTCCCAGATAAAAAGCTTGT
CAGATGGAGAAAGAAAACCTAGAAAAGCAATTAAAACAGATGTATTGCCACCC
TGTAAC TGCGGCTTGGTTTTCCACCTGCAAGATCCATGGGTACCAACAGGCCCT
GGAGCTGTGCAGAAGCAACGGGAGCACCCACCAGACTGTCAGTGGTATGCTCTT
GACCAGCTTCCGCCAGATGTACAACACAAGGCAAATGGTTTATCCTCAGTAAAG
AAAGTCCATCCGtaggcggccgc