

Division of Signal Transduction Therapy

Standard Operation Procedure

Preparation of Ubiquitin [K63wt]

| | |
|---|--|
| <u>Enzyme description:-</u> | Ubiquitin 1-76 [K63wt] |
| <u>Clone number:-</u> | DU24368 |
| <u>Source:-</u> | human recombinant |
| <u>Tag:-</u> | no tag |
| <u>Purification method:-</u> | Precipitation of contaminants, Source 15 S |
| <u>Expression system:-</u> | <i>E.coli</i> |
| <u>Calculated molecular mass:-</u> | |
| Monoisotopic | 8727 Da |
| Average Mass | 8732 Da |
| [cysteines reduced, methionines have not been oxidised] | |
| <u>Theoretical pI:-</u> | 7.7 |
| <u>Purity:-</u> | 95% |
| <u>Enzyme storage buffer:-</u> | H ₂ O |
| <u>Storage temperature:-</u> | -80°C |
| <u>Assay:-</u> | |

Division of Signal Transduction Therapy

Clone Data Sheet

Ubiquitin 1-76 [K63wt]

| | |
|---|--|
| <u>Protein</u> | Ubiquitin 1-76 [K63wt] |
| <u>Synonyms</u> | |
| <u>Clone Number</u> | DU24368 |
| <u>Species</u> | Human |
| <u>Accession Number</u> | Protein: P62987 |
| <u>Tags</u> | no tag |
| Aminoacid sequence of the expressed protein | MQIFVRTLTGRTITLEVEPSDTIENVRARIQDREGIPPDQQRLIFAGRQL EDGRTLSDYNIQKESTLHLVLRLRGG |
| Native sequence | In mammalian cells Ubiquitin is expressed as a precursor by several genes and cleaved by a DUB to become the mature 76 residue protein Ubiquitin. |
| Protease cleavage | N/A |
| Cloning sites | Nde1 / BamHI |
| <u>DNA sequence of insert</u> | CATATGCAAATCTCGTGAGGACCCCTGACTGGTAGGACCATCACTCTGAAG TGGAGCCGAGTGACACCATTGAGAATGTCAGGGCAAGGATCCAAGACAGGGA AGGCATCCCTCCTGACCAGCAGAGGTTGATCTTGCTGGGAGACAGCTGGAA GATGGACGCCACCCTGTCTGACTACAACATCCAGAAAGAGTCCACCCTGCACC TGGTCCTCCGTCTCAGAGGTGGGTGATAAAGATCT |