

Division of Signal Transduction Therapy

Standard Operation Procedure

Preparation of His-UBE2L6

<u>Enzyme description:-</u>	His-UBE2L6
<u>Clone number:-</u>	DU32148
<u>Source:-</u>	BL21 recombinant
<u>Tag:-</u>	N-terminal His ₆ -tag
<u>Purification method:-</u>	Ni ⁺⁺ -NTA-Sepharose
<u>Expression level:-</u>	5mg/L
<u>Calculated molecular mass:-</u>	
Monoisotopic	20200 Da
Average Mass	20212Da
[cysteines reduced, methionines have not been oxidised]	
<u>Theoretical pI:-</u>	7.7
<u>Purity:-</u>	90%
<u>Enzyme storage buffer:-</u>	
50mM HEPES pH 7.5, 150mM NaCl, 10% glycerol, 1mM DTT	
<u>Storage temperature:-</u>	-80°C

Assay:-

Loading assay with ISG15 and UBA7 in the presence of Mg-ATP

Division of Signal Transduction Therapy

Clone Data Sheet

His-UBE2L6

<u>Protein</u>	UBE2L6
<u>Synonyms</u>	UbcH6, RIG-B
<u>Clone Number</u>	DU32148
<u>Species</u>	Human
<u>Accession Number</u>	Protein: O14933 DNA: NM_004223
<u>Tags</u>	N-terminal His ₆ tag
Aminoacid sequence of the expressed protein	MGSSHHHHHHSSGLEVL <u>FQ</u> GPSMMASMRVVKELEDLQKKPPPYLRNLSS DDANVLVWHALLLPDQPPYHLKAFNLRISFPPEYPFKPPMIKF TT KIYHP NVDENGQICLP IISSENWKPC TK TCQVLEALNVLVNRPNIREPLRMDLAD LLTQNP EL FRKNAEEFTLRFGVDRPS
Native sequence	in bold
Protease cleavage	Prescission site underlined
Cloning sites	BamH1 / NotI
<u>DNA sequence of insert</u>	GGATCCATGATGGCGAGCATGCGAGTGGTGAAGGAGCTGGAGGATCTTCA GAAGAAGCCTCCCCCATACTGCGGAACCTGTCCAGCGATGATGCCAATG TCCTGGTGTGGCACGCTCTCCTCCTACCCGACCAACCTCCCTACCACCTG AAAGCCTTCAACCTGCGCATCAGCTTCCCGCCGGAGTATCCGTTCAAGCC TCCCATGATCAAATTCACAACCAAGATCTACCACCCCAACGTGGACGAGA ACGGACAGATTTGCCTGCCCATCATCAGCAGTGAGAACTGGAAGCCTTGC ACCAAGACTTGCCAAGTCCTGGAGGCCCTCAATGTGCTGGTGAATAGACC GAATATCAGGGAGCCCCTGCGGATGGACCTCGCTGACCTGCTGACACAGA ATCCGGAGCTGTTTCAGAAAGAATGCCGAAGAGTTCACCCTCCGATTCGGA GTGGACCGGCCCTCCTAAGCGGCCGC