Division of Signal Tranduction Therapy

Standard Operating Procedure

Preparation of active EPH A2 [591 – 976]

Enzyme description:- EPH A2 [591 - 976]

Clone number:- DU 4438

Source:- Recombinant

Expression system:- *E.coli*

Tag:- N-terminal GST

Purification method:- GSH Sepharose

Expression level:- 6 mg/L

Calculated molecular mass:-
Monoisotopic 70, 354.92 daltons
Average Mass 70, 400.53 daltons
[cysteines reduced, methionines have not been oxidised

Theoretical pI:- 6.94

Purity:- 90 %

Activation protocol:- Constitutively active

Enzyme storage buffer:-

50 mM Tris-HCl pH 7.5, 270 mM Sucrose, 150 mM NaCl, 0.1 mM EGTA,
0.1 % 2-mercaptoethanol, 0.02 % Brij-35, 1 mM benzamidine, 0.2 mM PMSF

Storage temperature:- -80 °C

Assay:- Standard filter binding assay

Assay buffer:-
50 mM Tris-HCl pH 7.5, 0.1 mM EGTA, 0.1 % 2–mercaptoethanol, 10 mM MgAc

Substrate:-
Poly Glu Tyr (4:1) Final concentration: 0.1 mg/ml

Specific activity range:- To be determined

Division of Signal Tranduction Therapy

Clone Data Sheet

EPH A2 [591 - 976]

Protein	EPH A2 [591 - 976]
Clone number	DU 4438
Species	Human
Accession number	NM_004431
Tags	N-terminal GST
Bacterially expressed protein	MSPILGYWKIKGLVQPTRLLLEYLEEKYEEHLYERDEGDKWRNKKFELG LEFPNLPYYIDGDVKLTQSMAIIRYIADKHNMLGGCPKERAEISMLEGA VLDIRYGVSRIAYSKDFETLKVDFLSKLPEMLKMFEDRLCHKTYLNGDH VTHPDFMLYDALDVVLYMDPMCLDAFPKLVCFKKRIEAIPQIDKYLKSS KYIAWPLQGWQATFGGGDHPPKSDLEVLFQGPLGS**PHTYEDPNQAVLKF** **TTEIHPSCVTRQKVIGAGEFGEVYKGMLKTSSGKKEVPVAIKTLKAGYT** **EKQRVDFLGEAGIMGQFSHHNIIRLEGVISKYKPMMIITEYMENGALDK** **FLREKDGEFSVLQLVGMLRGIAAGMKYLANMNYVHRDLAARNILVNSNL** **VCKVSDFGLSRVLEDDPEATYTTSGGKIPIRWTAPEAISYRKFTSASDV** **WSFGIVMWEVMTYGERPYWELSNHEVMKAINDGFRLPTPMDCPSAIYQL** **MMQCWQQERARRPKFADIVSILDKLIRAPDSLKTLADFDPRVSIRLPST** **SGSEGVPFRTVSEWLESIKMQQYTEHFMAAGYTAIEKVVQMTNDDIKRI** **GVRLPGHQKRIAYSLLGLKDQVNTVGIPI**
Native sequence	Amino acids P591 – I976 (end) of human EPH A2. Residue P232 of the fusion protein is equivalent to P591 of the native enzyme. The GST tag is located at residues 1 – 220.
Protease cleavage	PreScission (LEVLFQGPL) residues 221 - 229
Cloning sites	*Bgl*III and *Sal*1site of pGEX 6P-1

Nucleotide Sequence

atgtcccctatactaggttattggaaaattaagggccttgtgcaaccca
ctcgacttcttttggaatatcttgaagaaaaatatgaagagcatttgta
tgagcgcgatgaaggtgataaatggcgaaacaaaaagtttgaattgggt
ttggagtttcccaatcttccttattatattgatggtgatgttaaattaa
cacagtctatggccatcatacgttatatagctgacaagcacaacatgtt
gggtggttgtccaaaagagcgtgcagagatttcaatgcttgaaggagcg
gttttggatattagatacggtgtttcgagaattgcatatagtaaagact
ttgaaactctcaaagttgattttcttagcaagctacctgaaatgctgaa
aatgttcgaagatcgtttatgtcataaaacatatttaaatggtgatcat
gtaacccatcctgacttcatgttgtatgacgctcttgatgttgttttat
acatggacccaatgtgcctggatgcgttcccaaaattagtttgttttaa
aaaacgtattgaagctatcccacaaattgataagtacttgaaatccagc
aagtatatagcatggcctttgcagggctggcaagccacgtttggtggtg
gcgaccatcctccaaaatcggatctggaagttctgttccaggggcccct
gggatctCCCCACACATATGAGGACCCCAACCAGGCTGTGTTGAAGTTC
ACTACCGAGATCCATCCATCCTGTGTCACTCGGCAGAAGGTGATCGGAG
CAGGAGAGTTTGGGGAGGTGTACAAGGGCATGCTGAAGACATCCTCGGG
GAAGAAGGAGGTGCCGGTGGCCATCAAGACGCTGAAAGCCGGCTACACA
GAGAAGCAGCGAGTGGACTTCCTCGGCGAGGCCGGCATCATGGGCCAGT
TCAGCCACCACAACATCATCCGCCTAGAGGGCGTCATCTCCAAATACAA
GCCCATGATGATCATCACTGAGTACATGGAGAATGGGGCCCTGGACAAG
TTCCTTCGGGAGAAGGATGGCGAGTTCAGCGTGCTGCAGCTGGTGGGCA
TGCTGCGGGGCATCGCAGCTGGCATGAAGTACCTGGCCAACATGAACTA
TGTGCACCGTGACCTGGCTGCCCGCAACATCCTCGTCAACAGCAACCTG
GTCTGCAAGGTGTCTGACTTTGGCCTGTCCCGCGTGCTGGAGGACGACC
CCGAGGCCACCTACACCACCAGTGGCGGCAAGATCCCCATCCGCTGGAC
CGCCCCGGAGGCCATTTCCTACCGGAAGTTCACCTCTGCCAGCGACGTG
TGGAGCTTTGGCATTGTCATGTGGGAGGTGATGACCTATGGCGAGCGGC
CCTACTGGGAGTTGTCCAACCACGAGGTGATGAAAGCCATCAATGATGG
CTTCCGGCTCCCCACACCCATGGACTGCCCCTCCGCCATCTACCAGCTC
ATGATGCAGTGCTGGCAGCAGGAGCGTGCCCGCCGCCCCAAGTTCGCTG
ACATCGTCAGCATCCTGGACAAGCTCATTCGTGCCCCTGACTCCCTCAA
GACCCTGGCTGACTTTGACCCCCGCGTGTCTATCCGGCTCCCCAGCACG
AGCGGCTCGGAGGGGGTGCCCTTCCGCACGGTGTCCGAGTGGCTGGAGT
CCATCAAGATGCAGCAGTATACGGAGCACTTCATGGCGGCCGGCTACAC
TGCCATCGAGAAGGTGGTGCAGATGACCAACGACGACATCAAGAGGATT
GGGGTGCGGCTGCCCGGCCACCAGAAGCGCATCGCCTACAGCCTGCTGG
GACTCAAGGACCAGGTGAACACTGTGGGGATCCCCATCtgagtcgac